

Please mark only **ONE** BOX with a **X**

When should you re-evaluate your management of a foot ulcer that is not infected or healing?	<input type="checkbox"/> 1 week <input type="checkbox"/> 2 weeks <input type="checkbox"/> 1 month <input type="checkbox"/> 2 months <input type="checkbox"/> 4 months <input type="checkbox"/> Sundays
How do you most definitively clinically determine if an ischaemic ulcer requires a Vascular Surgical referral?	<input type="checkbox"/> Look at skin <input type="checkbox"/> Ask patient <input type="checkbox"/> Palpate pulses <input type="checkbox"/> Toe pressures <input type="checkbox"/> Pole test
If your patient has >2cm Cellulitis around their foot ulcer what would you do?	<input type="checkbox"/> Initiate antibiotics <input type="checkbox"/> Panic <input type="checkbox"/> Get MRI <input type="checkbox"/> Refer for potential admission <input type="checkbox"/> Dress with honey
What is the most appropriate Wound Dressing for non-infected plantar Neuropathic Ulcers?	<input type="checkbox"/> Hydrocolloid <input type="checkbox"/> Foam <input type="checkbox"/> Film <input type="checkbox"/> Honey <input type="checkbox"/> Hydrofibre <input type="checkbox"/> Combine
How do you measure the size of a foot ulcer according to the International group?	<input type="checkbox"/> Draw picture <input type="checkbox"/> Take photo <input type="checkbox"/> Measure length & width (mm ²) <input type="checkbox"/> Pray <input type="checkbox"/> Ask patient
How do you primarily assess the depth of a foot ulcer?	<input type="checkbox"/> Look <input type="checkbox"/> Fingers <input type="checkbox"/> X-ray <input type="checkbox"/> MRI <input type="checkbox"/> Sterile probe <input type="checkbox"/> Refer Endocrinologist
What is the most important principle to manage a non-infected Neuropathic Ulcer?	<input type="checkbox"/> Wound dressing <input type="checkbox"/> Antibiotics <input type="checkbox"/> Off-load pressure <input type="checkbox"/> Refer Vascular Surgeon